

AFTER CARE INSTRUCTIONS: FILLER

IMMEDIATELY AFTER TREATMENT

Redness and swelling are normal.

Bruising may also be visible.

You may experience some tenderness at the treatment site(s) that can last for a few hours or a few days.

Depending upon the area(s) treated and product(s) used, you may feel "firmness" in the treated area(s) for 1 to 2 weeks after treatment.

Over time, the area(s) will soften and "settle"

POST-TREATMENT INSTRUCTIONS

Apply an ice or cold gel pack to the area(s) treated (avoiding pressure) as this helps reduce swelling and the potential for bruising.

Once you have adequately cooled/iced the area(s) as instructed and any pinpoint bleeding from the injection site(s) has subsided, you may begin wearing makeup.

Avoid placing excessive pressure on the treated area(s) for the first few hours and up to 2 to 3 days; when cleansing your face or applying makeup, be very gentle.

Avoid exercise or strenuous activities for the remainder of the treatment day; you may resume other normal activities/routines after three days

You may take Acetaminophen/ Tylenol if you experience any mild tenderness or discomfort.

Avoid extended UV exposure until any redness/swelling has subsided. Be sure to apply an SPF 30 or higher sunscreen

Wait a minimum of four weeks (or as directed by your provider) before receiving any skin care or laser treatments.

PLEASE CALL US WITH ANY QUESTIONS OR CONCERNS
248-413-5967

FOR AFTER HOURS QUESTIONS AND EMERGENCIES,
Please Contact Roxanne at 989-770-8110